

Light soils reflect more heat than darker soils which are better absorbers. Such soil differences may give rise to slight variations in the temperature of the region.

As a whole, dry soils like sands are very sensitive to temperature changes, whereas wet soils, like clay, retain much moisture and warm up or cool down more slowly.

PRECIPITATION

Types of Precipitation.

If air is sufficiently cooled below dew-point, tiny drops of water vapour will condense around dust particles. When they float about as masses of minute water droplets or ice crystals at a considerable height above sea level, they form clouds—cirrus, cumulus or stratus. When condensation occurs at ground level without necessarily resulting in rain, haze, mist or fogs are formed. In higher latitudes or altitudes, where condensation of water vapour may take place in the atmosphere at temperatures below freezing-point, snow falls, either as feathery flakes or individual ice crystals. If the moist air ascends rapidly to the cooler layers of the atmosphere, the water droplets freeze into ice pellets and fall to the earth as hail or hailstones.

As more and more super-cooled water drops accumulate around a hailstone, it increases steadily in size; some of them weigh as much as two pounds. In a severe hail-storm the hailstones do great damage to crops and buildings. Very often, the ice-pellets exist as frozen rain-drops, melting and re-freezing on their way down; this forms sleet. It is only when the droplets in clouds coalesce into larger drops between 0.2mm and 6mm that rain falls.

RAINFALL

Types of Rainfall.

There are three major types of rainfall.

1. Convictional rainfall.

This type of rainfall is most common in regions that are intensely heated, either during the day, as in the tropics, or in the summer, as in temperate interiors. When the earth's surface is heated by conduction, moisture-laden vapour rises because heated air always expands, and becomes lighter. Air rises in a convection current after a prolonged period of intense heating. In ascending, its water vapour condenses into cumulonimbus clouds with a great vertical extent.

This probably reaches its maximum in the afternoon when the convectional system is well developed. Hot, rising air has great capacity for holding moisture, which is abundant in regions of high relative humidity. As the air rises it cools and when saturation point is reached torrential downpours occur, often accompanied by thunder and lightning. The summer showers in temperate regions are equally heavy with occasional thunderstorms. These downpours may not be entirely useful for agriculture because the rain is so intense that it does not sink into the soil but is drained off almost immediately.

2. Orographic or relief rain.

Unlike convectional rain which is caused by convection currents, Orographic rain is formed wherever moist air is forced to ascend a mountain barrier. It is best developed on the windward slopes of mountains where the prevailing moisture-laden winds come from the sea. The air is compelled to rise, and is thereby cooled by expansion in the higher altitudes and the subsequent decrease in atmospheric pressure.

Further ascent cools the air until the air is completely saturated (relative humidity is 100 percent). Condensation takes place forming clouds and eventually rain. Since it is caused by the relief of the land, it is also known as relief rain. Much of the precipitation experienced on the windward slopes of the north-east of West Malaysia, western New Zealand, western New Zealand, western Scotland and Wales and the Assam hills of the Indian sub-continent, is relief rain.

On descending the leeward slope, a decrease in altitude increases both the pressure and the temperature; the air is compressed and warmed. Consequently, the relative humidity will drop. There is evaporation and little or no precipitation. The area in the lee of the hills is termed the rain shadow area. The effects of rain shadow are felt on the Canterbury Plain of South Island, New Zealand and the western slopes of the Northern and Central Andes and in many other areas.

3. Cyclonic or frontal rain.

This type of rainfall is independent of relief or convection. It is purely associated with cyclonic activity whether in the temperate regions (depressions) or tropical regions (cyclones). Basically it is due to the convergence (meeting) of two different air masses with different temperatures and other physical properties. As cold air is denser, it tends to remain close to the ground. The warm air is lighter and tends to rise over the cold air. In ascent pressure decreases, the air expands and cools, condensation takes place and light showers called cyclonic or frontal rain occur. The heavier and colder air masses eventually push up the warmer and lighter air and the sky is clear again.

PRESSURE AND PLANETARY WINDS

World pressure belts.

The circulation of water in the oceans and noted that they follow a regular pattern, flowing from the poles equatorwards and from the equator polewards. In the same way, there is also a circulation of air over the surface of the earth caused by the differences in pressure.

Along the equator and within 5 degrees north and south, is the Equatorial Low Pressure Belt, where there is intense heating, with expanding air and ascending convection currents. This equatorial belt is often termed the Doldrums, because sailors in the olden days often found themselves becalmed here. It is a zone of wind convergence.

About 30° N. and S. are two Temperate Low Pressure Belts which are also zones of convergence with cyclonic activity. The sub-polar low pressure areas are best developed over the oceans, where temperature difference between summer and winter are negligible.

At The North and South Poles 90° N and S. where temperatures are permanently low, are the Polar High Pressure Belts. Unlike the water masses of the high latitudes in the southern hemisphere, high pressures of the corresponding latitudes in the northern hemisphere are a little complicated by the presence of much land. Some pressure differences between summer and winter can be expected.

The Planetary winds.

Within this pattern of permanent pressure belt on the globe, winds tend to blow from the high pressure belts to the low pressure belts as the planetary winds.

Instead of blowing directly from one pressure belt to another, however, the effect of the rotation of the earth (Coriolis Force) tends to deflect the direction of the winds. In the northern hemisphere, winds are deflected to their right, and in the southern hemisphere to their left. This is known as Ferrel's Law of Deflection. The Coriolis force is absent along the equator but increased progressively towards the poles.

For this reason, winds blowing out from the Sub-Tropical High Pressure Belt in the northern hemisphere towards the Equatorial Low become North-East Trade Winds and those in the southern hemisphere become the South-East Trade winds. These trade winds are the most regular of all the planetary winds. They blow with great force and in a constant direction. They were thus helpful to early traders who depended on the wind when sailing the high seas; hence the name 'trade winds'. Since they blow from the cooler sub-tropical latitudes to the warmer tropics, they have great capacity for holding moisture. In their passage across the open oceans, they gather more moisture and bring heavy rainfall to the east coasts of continents within the tropics. As they are off-shore on the west coast, these regions suffer from great aridity and form the Trade Wind Hot Deserts of the world, e.g. the Sahara, Kalahari Atacama and Great Australian Deserts.

From the Sub-Tropical High Pressure Belts, winds blow towards the Temperate Low Pressure Belts as the variable Westerlies. Under the effect of the Coriolis force, they become the South-Westerlies in the northern hemisphere and the North-Westerlies in the southern hemisphere. They are more variable in the northern hemisphere, but they play a valuable role in carrying warm equatorial waters and winds to western coasts of temperate lands.

This warming effect and other local pressure differences have resulted in a very variable climate in the temperate zones, dominated by the movements of cyclones and anticyclones. In the southern hemisphere where there is a large expanse of ocean, from 40° S to 60° S., Westerlies blow with much greater force and regularity throughout the year. They bring much precipitation to the western coasts of continents. The weather is damp and cloudy and the seas are violent and stormy. It is thus usual for seafarers to refer to the Westerlies as the Roaring Forties, Furious Fifties and Shrieking or Stormy Sixties, according to the varying degree of storminess in the latitudes in which they blow.

It must be pointed out that not all the western coasts of the temperate zone receive Westrlies throughout the year. Some of them like California, Ibria, central Chile, southern Africa and south Western Australia receive Westerlies only in winter. This is caused by the 'shifting of the wind belts' of such regions which lie approximately between the latitudes 30° and 40° N and S. Due to the earth's inclination, the sun is overhead at midday in different parts of the earth at different seasons. The entire system of pressure and wind belts follows he movement of the midday and wind belts follows the movement of the midday sun. In June when the overhead sun is over the Tropic of Cancer, all the belts move about 5° - 10° south of their average position. The 'Mediterranean' parts of Europe and California then come under the influence of the Westerlies and receive rain in December (winter in the northern hemisphere).

Lastly, mention must be made of the Polar Easterlies which blow out from the Polar High Pressure Belts towards the Temperature Low Pressure Belts. These are extremely cold winds as they come from the tundra and ice-cap regions. They are more regular in the south than in the north.

LAND AND SEA BREEZES AND MONSOONS

Land and sea breezes are, in fact, monsoons on a smaller scale. Both are basically caused by differential heating of land and sea, the former in a diurnal rhythm and the latter in a seasonal rhythm.

During the day, the land gets heated up much faster than the sea. Warm air rises forming a region of local low pressure. The sea remains comparatively cool with a higher pressure so a sea breeze blows in from sea to land. Its speed or strength is between 5-20m.p.h and it is generally stronger in tropical than temperate regions. Its influence does not normally exceed 15miles from the coast. It is most deeply felt when one stands facing the sea in a coastal resort.

At night the reverse takes place. As the land cools down much faster than the sea, the cold and heavy air produces a region of local high pressure. The sea conserves its heat and remains quite warm. Its pressure is comparatively low. A land breeze thus blows out from land to sea. Fishermen in the tropics often take advantage of the out-going land breeze and sail out with it. They return the next morning with the in-coming sea breeze, complete with their catch.

In the same way, monsoons are caused. Rapid heating in the hot summer over most parts of India for example induces heated air to rise. The South-West Monsoon from the surrounding ocean is attracted by the low pressure over the land and blows in, bringing torrential rain to the sub-continent.

Similarly, in winter when the land is cold, the surrounding seas remain comparatively warm. High pressure is created over Indo-Pakistan and the North-East Monsoon blows out from the continent into the Indian Ocean and the Bay of Bengal.

FOHN WIND OR CHINOOK WIND

Both the Fohn and Chinook winds are dry winds experienced on the leeward side of mountains when experienced on the leeward side of mountains when descending air becomes compressed with increased pressure. The Fohn win is experienced in the valleys of the northern Alps, particularly in Switzerland in spring. Chinook winds are experienced on the eastern slopes on the eastern slopes of the Rockies in U.S.A. and Canada in winter.

Air ascending the southern slopes of the Alps expands and cools. Condensation takes place when the air is saturated. Rain and even snow fall on the higher slopes.

In descending the northern slope, the wind experiences and increase in pressure and temperature. The air is compressed and warmed. Most of its moisture is lost and the wind reaches the valley bottom as a dry, hot wind – the Fohn. It may raise the temperature by 15° to 30°F., within an hour! It melts snow and causes avalanches. In North America it is called Chinook, meaning ‘the snow-eater’. But it has its blessings too, it hastens the growth of crops and fruits and thaws the snow-covered pastures. In the Rockies, the Chinook has been known to raise temperature 35°F. within 15 minutes! The occurrence of frequent Chinooks means winter is mild.

CYCLONIC ACTIVITY

Tropical cyclones, typhoons, hurricanes and tornadoes

All These are different kinds of tropical cyclones. They are well developed low pressure systems into which violent winds blow. Typhoons occur in the China Sea; tropical cyclones in the Indian Ocean; hurricanes in the West Indian islands in the Caribbean; tornadoes in the Guinea lands of West Africa, and the southern U.S.A. in which the local name of Whirl-wind is often applied and willy-willies occur in north-western Australia.

Typhoons occur mainly in regions between 6° and 20° north and south of the equator and are most frequent from July to October. In extent, they are smaller than temperate cyclones and have a diameter of only 50 to 200 miles, but they have a much steeper pressure gradient. Violent winds with a velocity of over 100 m.p.h are common. The sky is overcast and the torrential downpour is accompanied by thunder and lightning. In the wake of the typhoon, damage is widespread, e.g. in 1922, a typhoon that hurled huge waves on to the Swatow coast drowned 50,000 people.

The other tropical cyclones have similar characteristics and differ, perhaps, only in intensity, duration and locality. Hurricanes have calm, rainless centres where the pressure is lowest (about 965 mb.) but around this 'eye' the wind strength exceeds force 12 of the Beaufort scale (75 m.p.h). Dense dark clouds gather and violent stormy weather lasts for several hours. A terrible hurricane struck Barbados in the West Indies in 1780, which nearly destroyed the whole island, tearing down buildings and uprooting trees. About 6,000 inhabitants were reported dead.

Tornadoes are small but very violent tropical and sub-tropical cyclones in which the air is spiraling at a tremendous speed of as much as 500 m.p.h! A tornado appears as a dark funnel cloud 250 to 1,400 feet in diameter. As a tornado passes through a region, it writhes and twists, causing complete devastation within the limits of its passage. There is such a great difference in pressure that houses virtually explode. Tornadoes are most frequent in spring but not common in many countries and their destructive effects are confined to a small area. Tornadoes are most typical of the U.S.A and occur mainly in the Mississippi basin.

Cyclones.

These are better known as depressions and are confined to temperate latitudes. The lowest pressure is in the centre and the isobars, as shown in climatic charts, are close together. Depressions vary from 150 to 2,000 miles in extent.

They remain quite stationary or move several hundred miles in a day. The approach of a cyclone is characterized by a fall in barometric reading, dull sky, Oppressive air and strong winds. Rain or snow falls and the weather is generally bad. Winds blow inwards into regions of low pressure in the centre, circulating in anticlockwise direction in the northern hemisphere and clockwise in the southern hemisphere . Precipitation resulting from cyclonic activities is due to the convergence of warm tropical air and cold polar air. Fronts are developed and condensation takes place, forming either rain , snow or sleet.

Anticyclones.

These are the opposite of cyclones, with high pressure in the centre and the isobars far apart. The pressure gradient is gentle and winds are light. Anticyclones normally herald fine weather. Skies are clear, the air is calm and temperatures are high in summer but cold in winter. In winter intense cooling of the lower atmosphere may result in thick fogs. Anticyclonic conditions may last for days or weeks and then fade out quietly. Winds in anticyclones blow outwards and are also subject to deflection, but they blow clockwise in the northern hemisphere and anticlockwise in the southern hemisphere.

CLIMATIC TYPES AND NATURAL VEGETATION

It is necessary to divide the world into several climatic zones, each with its own climatic characteristics, natural vegetation (forests, grasslands or deserts), crops, animals and human activities. Though the geographical characteristics may not be absolutely uniform in each climatic type, they have many things in common. The scheme of the world's climatic types with their seasonal rainfall and natural vegetation.

<u>Climatic Zone</u>	<u>Latitude</u>	<u>Climatic Type</u>	<u>Rainfall Regime</u>	<u>Natural Vegetation</u>
----------------------	-----------------	----------------------	------------------------	---------------------------

1. Equatorial Zone	0° - 10° N & S	1. Hot, wet equatorial	Rainfall all year round: 80 inches	Equatorial rain forests
2. Hot Zone	10° - 30° N & S	2. (a) Tropical Monsoon	Heavy summer rain: 60 inches	Monsoon forests
		(b) Tropical Marine	Much summer rain: 70 inches	
		3. Sudan Type	Rain mainly in Summer: 30 inches	Savanna (tropical grassland)
		4. Desert : (a) Saharan type (b) Midlatitude	Little rain: 5 inches	Desert vegetation and scrub
3. Warm Temperate Zone	30° - 45° N & S	5. Western Margin (Mediterranean type)	Winter rain: 35 inches	Mediterranean forests and shrub
		6. Central Continental (steppe type)	Light summer rain: 20 inches	Steppe or temperate grassland
		7. Eastern Margin: (a) China type (b) Gulf type (c) Natal type	Heavier summer rain: 45 inches	Warm, wet forest and bamboo
4. Cool Temperate Zone	45° - 65° N & S	8. Western Margin (British type)	More rain in autumn and winter: 30 inches	Deciduous forests
		9. Central continental (Siberian type)	Light summer rain: 25 inches	Evergreen coniferous forests
		10. Eastern Margin (Laurentian type)	Moderate summer rain: 40 inches	Mixed forest (coniferous and deciduous)
5. Cold Zone	65° - 90° N & S	11. Arctic or Polar	Very light summer rain: 10 inches	Tundra, mosses, lichens
6. Alpine Zone		12. Mountain climate	Heavy rainfall (variable)	Alpine pasture, conifers, fern, snow.